

Closedness in the Weak Topology of the Dual Pair L^1, \mathcal{C}

MICHEL VALADIER

*Université II, 34060 Montpellier Cedex, France**Submitted by J. L. Lions*

This paper gives improvements and new results along the lines of the work of Rockafellar, Olech, and Sainte-Beuve. It is concerned with closure of sets and lower semicontinuity of functionals relative to the weak topologies on L^1 or on the space \mathcal{M} of measures.

1. INTRODUCTION

This paper is in the line of Rockafellar [11], Olech [6, 7], and Sainte-Beuve [13, 17].

Let T be a topological space, μ a positive measure on T , and E a separable Fréchet space. Let Γ be a measurable multifunction from T to E' , the dual space of E . We denote by \mathcal{L}_T^1 the set of μ -integrable selections of Γ , by L_T^1 its quotient. Let \mathcal{C}_E^b be a vector space of E -valued continuous bounded functions on T . Consider the geometric problems: What is the closure of L_T^1 for the weak topology $\sigma(L_E^1, \mathcal{C}_E^b)$, what is the weak closure of L_T^1 in the space $\mathcal{M}_{E'}$ of E' -valued measure? Now let $f: T \times E' \rightarrow \mathbf{R}$ be a measurable function and define $I_f(u)$ for $u \in L_E^1$, by $I_f(u) = \int f(t, u(t)) \mu(dt)$. Consider the functional problems: What is the weak l.s.c. hull of I_f , what is the polar (calculated on $\mathcal{M}_{E'}$) of its polar?

First Rockafellar in [11] studied the functional problem and gave corollaries about the geometric problem. His work is an application of his results on the duality between L^∞ and $L^{\infty'}$. Then Olech [6] treated the geometric problem by a completely different approach. In [7] he used [6] to prove that I_f is l.s.c. iff L_T^1 is closed (where $\Gamma(t)$ is the epigraph of $f(t, \cdot)$). Sainte-Beuve in [13] weakens the hypotheses of [6]: T becomes a completely regular Suslin space instead of compact metric, and E becomes a separable Fréchet space instead of \mathbf{R}^n . In [17] Sainte-Beuve extends [7].

Applications of those kinds of results are given in Olech [8, 9]. Some other similar problems have been recently studied in Brézis [1], Ioffe [3, 4], and Tran cao Nguyen [20].

In Section 2 we set notation and hypotheses. In Section 3 we state briefly the foregoing results we need. In Section 4 we give some useful examples. In

Section 5 we give our results about the geometric problem. We extend a necessary and sufficient condition for L_T^1 to be closed (Olech [6, Corollary 2, p. 317]) within the framework of Sainte-Beuve. Then we give a new formula for the closure of L_T^1 . In Section 6 we apply our geometric results to characterize the bipolar of I_f . The formula is similar to [11, p. 458]. The main improvements are: (1) Rockafellar needs hypotheses which ensure that I_f is l.s.c., (2) he assumes $E = \mathbf{R}^n$, and (3) he deduces the formula from a rather technical preliminary result [11, Theorem 4, p. 452].

Nevertheless in [11] T is not assumed to be separable. Then we give necessary and sufficient conditions for I_f to be l.s.c. and a description of its l.s.c. hull.

All the results of Section 5 (except 3 of Corollary 4) are valid when T is a completely regular Suslin space and μ is σ -finite. But, in Section 6 we suppose that T is compact. To avoid changing the hypotheses and notation in the middle of the text, we assume throughout the paper that T is compact. One can find in [22] the general version; the proofs are identical. Finally we wish to thank M. F. Nougues-Sainte-Beuve for a very helpful conversation.

2. NOTATIONS AND HYPOTHESES

[Let T be a compact metric space, μ a positive bounded measure on T , and E a separable Fréchet space. We denote by $\mathcal{C}_E(T)$ (or shortly \mathcal{C}_E , even \mathcal{C} when $E = \mathbf{R}$) the space of continuous E -valued functions on T . When it is endowed with the uniform topology its dual identifies with a subset of the set of E' -valued measures.

We denote by $\mathcal{B}(E'_s)$ the Borel tribe of the $\sigma(E', E)$ topology, by $\mathcal{B}(T)$ the Borel tribe of T , and by $\mathcal{B}(T)_\mu$ its μ -completion. It is well known that E'_s is a Lusin (or standard) space, hence, a Suslin space, so that one can define μ -measurability of a function $u: T \rightarrow E'_s$ in any of the classical ways (see [2, Theorem III.36, p. 83]).

For any positive measure θ on T we denote by $\mathcal{L}_{E'}^1([E], \theta)$ the space of all θ -measurable functions $u: T \rightarrow E'_s$ such that there exist $\alpha \in \mathcal{L}_{\mathbf{R}}^1(\theta)$ and an equicontinuous set $Q \subset E'$ such that $u(t) \in \alpha(t)Q$ θ -a.e. And $L_{E'}^1([E], \theta)$ denotes the quotient space. We shall write $\mathcal{L}_{E'}^1$ and $L_{E'}^1$ for $\mathcal{L}_{E'}^1([E], \mu)$ and $L_{E'}^1([E], \mu)$.

For every θ , and every $u \in \mathcal{L}_{E'}^1([E], \theta)$ the function u is θ -Pettis integrable, so that the formula

$$m(A) = \int_A u \, d\theta$$

defines an E' -valued measure. It is denoted by $u \cdot \theta$. The set of all measures $u \cdot \theta$ for $u \in \mathcal{L}_{E'}^1([E], \theta)$ is by definition [13] the set of E' -valued measures with bounded variation and absolutely continuous with respect to θ . The space of all vector-valued measures with bounded variation (for all θ) is denoted by $\mathcal{M}_{E'}(T)$

or shortly $\mathcal{M}_{E'}$. It is the dual space of \mathcal{C}_E . So we have at least three pairs of spaces in separated duality: E and E' , \mathcal{C}_E and $L_{E'}^1$, \mathcal{C}_E and $\mathcal{M}_{E'}$. Let C be a subset of one of these spaces, then $\delta^*(\cdot | C)$ denotes its support function and C^0 its polar: for example, if $C \subset E'$

$$\delta^*(x | C) = \sup\{\langle x', x \rangle | x' \in C\} \quad \text{for } x \in E,$$

$C^0 = \{x \in E | \forall x' \in C \langle x', x \rangle \leq 1\}$. And $\text{co } C$ denotes the convex hull of C . If C is closed and convex, then $\text{As } C$ denotes the asymptotic cone of C [2, I.7, p. 8], and C is said to be LCNL if it is weakly locally compact and contains no line [2, I.15, p. 16–17].

Let Γ be a multifunction from T to E' . We say that it is μ -measurable if its graph belongs to $\mathcal{B}(T)_\mu \otimes \mathcal{B}(E')$.

We denote by \mathcal{L}_E^1 the set $\{u \in \mathcal{L}_{E'}^1 | u(t) \in \Gamma(t) \text{ } \mu \text{ a.e.}\}$ and by L_Γ^1 its quotient. We denote by $\overline{L_\Gamma^1}$ the $\sigma(L_{E'}^1, \mathcal{C}_E)$ closure of L_Γ^1 and by $\overline{\overline{L_\Gamma^1}}$ its $\sigma(\mathcal{M}_{E'}, \mathcal{C}_E)$ closure. We shall denote by \mathcal{H} the conic convex set $\text{dom } \delta^*(\cdot | L_\Gamma^1) = \{\varphi \in \mathcal{C}_E | \delta^*(\varphi | L_\Gamma^1) < \infty\}$.

Hypotheses

(H1) μ is nonatomic or $\Gamma(t)$ is μ a.e. convex. (This hypothesis could be weakened as in [12]: Γ is supposed convex valued on every atom of μ .)

(H2) L_Γ^1 is nonempty.

3. FOREGOING RESULTS

The set $\overline{\overline{L_\Gamma^1}}$ has two important properties: (1) It is convex, thanks to the Liapunov theorem, (2) it is the intersection of a countable family of closed half spaces. This follows from the fact that $\mathcal{M}_{E'}$ has the Lindelöf property. (When E is Fréchet and T completely regular Suslin that result has been proved by Sainte-Beuve.) So there exists a sequence (φ_n) in \mathcal{C}_E such that $\forall n$, $\delta^*(\varphi_n | L_\Gamma^1) < \infty$ and

$$\overline{\overline{L_\Gamma^1}} = \{m \in \mathcal{M}_{E'} | \forall n, \langle m, \varphi_n \rangle \leq \delta^*(\varphi_n | L_\Gamma^1)\}.$$

We can now state the Olech theorem extended by Sainte-Beuve:

THEOREM 0. *Putting*

$$\Sigma(t) = \{x' | \forall n, \langle x', \varphi_n(t) \rangle \leq \delta^*(\varphi_n(t) | \Gamma(t))\}$$

$$C(t) = \{x' | \forall n, \langle x', \varphi_n(t) \rangle \leq 0\},$$

one has

$$\overline{\overline{L_\Gamma^1}} = L_E^1$$

and

$$\overline{L_E^1} = L_{\Sigma^1} + \{m \mid \exists \theta, \exists u \in L_{C^1}(\theta) \text{ such that } m \sim u \cdot \theta\}.$$

In the last formula one can consider only measures θ which are μ -singular. Moreover $\forall t, C(t) = [\mathcal{H}(t)]^0$ (where $\mathcal{H}(t) = \{\varphi(t) \mid \varphi \in \mathcal{H}\}$).

Remark 1. It is easy to see that $\mathcal{H}(t)$ and $C(t)$ depend only on the values of Γ on a neighborhood of t . Indeed: $\mathcal{H}(t) = \{\varphi(t) \mid \varphi \in \mathcal{C}_E, \exists V \text{ open such that } t \in V \text{ and } \int_V \delta^*(\varphi(s) \mid \Gamma(s)) \mu(ds) < \infty\}$, because if φ belongs to the right-hand side, then there exists a continuous function $\alpha: T \rightarrow [0, 1]$ such that $\alpha(t) = 1$ and $\alpha(s) = 0$ if $s \notin V$. Then $\alpha\varphi \in \mathcal{H}$ and $\varphi(t) = (\alpha\varphi)(t)$ belongs to $\mathcal{H}(t)$.

Remark 2. It is obvious from the definition of Σ and C that $\Sigma(t) \neq \phi$ and $\delta^*(\varphi_n(t) \mid \Gamma(t)) < \infty$ implies $C(t) = \text{As } \Sigma(t)$. As L_{Γ^1} is nonempty, L_{Σ^1} is also nonempty and so $\Sigma(t) \neq \phi$ a.e. Hence one has $C(t) = \text{As } \Sigma(t)$ a.e.

THEOREM 1. *The set L_{Γ^1} is $\sigma(L_{E^1}^1, \mathcal{C}_E)$ closed iff there exist two sequences (φ_n) in \mathcal{C}_E and (ψ_n) in $\mathcal{L}_{\mathbb{R}}^1$ such that*

$$\Gamma(t) = \{x' \mid \forall n, \langle x', \varphi_n(t) \rangle \leq \psi_n(t)\} \quad \mu \text{ a.e.} \quad (1)$$

Remark 3. The above necessary and sufficient condition is not very practical. Olech gave also another one, which we extend in Theorem 3.

Remark 4. The relation $\psi_n \in \mathcal{L}_{\mathbb{R}}^1$ is essential in the proof of sufficiency: see Example 1 below.

4. EXAMPLES

EXAMPLE 1. Let $T = [0, 1]$, $\mu = dt + \delta_0$ where dt is the Lebesgue measure and δ_0 the Dirac measure, $E = \mathbf{R}$ and

$$\begin{aligned} \Gamma(t) &= \{0\} & \text{if } t &= 0 \\ &= [0, 1/t] & \text{if } t > 0. \end{aligned}$$

Then for $n \geq 1$ the function $u_n = n\chi_{[0, 1/n]}$ belongs to $\mathcal{L}_{\Gamma^1}^1$, but it converges weakly to $\chi_{\{0\}}$ which does not belong to $\mathcal{L}_{\Gamma^1}^1$. It will be seen after Theorem 3 that $\overline{L_{\Gamma^1}^1} = L_{\Sigma^1}$ with

$$\begin{aligned} \Sigma(t) &= [0, \infty[& \text{if } t &= 0 \\ &= [0, 1/t] & \text{if } t > 0. \end{aligned}$$

Finally we notice that (1) is satisfied with a nonintegrable ψ_n :

$$\Gamma(t) = \{x' \mid \langle x', \varphi_1(t) \rangle \leq \psi_1(t) \text{ and } \langle x', \varphi_2(t) \rangle \leq \psi_2(t)\}$$

with

$$\begin{aligned} \varphi_1 &= -1, & \psi_1 &= 0, & \varphi_2 &= 1, \\ \psi_2(t) &= 0 & \text{if } t &= 0 \\ &= 1/t & \text{if } t &> 0. \end{aligned}$$

EXAMPLE 1'. In Example 1 μ has an atom. We can introduce an auxiliary variable: Let $T = [0, 1]^2$, $\mu = (dt + \delta_0) \otimes ds$,

$$\begin{aligned} \Gamma(t, s) &= \{0\} & \text{if } t &= 0 \\ &= [0, 1/t] & \text{if } t > 0. \end{aligned}$$

Thus μ is nonatomic but the same phenomena happens.

EXAMPLE 2. Let $T = [0, 1]$, $\mu = dt$, $E = \mathbf{R}$, and let K be an analogous of the Cantor set but with positive measure. Let

$$\begin{aligned} \Gamma(t) &= \{0\} & \text{if } t \in K \\ &= [0, \infty[& \text{if } t \in T - K. \end{aligned}$$

It will be proved after Theorem 3 that $\overline{L_R^1} = L_{[0, \infty[}^1$. (A similar example is given in [8, p. 622].)

Remark 5. One could think that a necessary condition (which would fail in the examples) for L_R^1 to be closed is a kind of semicontinuity of Γ . This is not true in an elementary way: Indeed if a, b belong to $\mathcal{L}_{\mathbf{R}}^1$ with $a \leq b$, and $\Gamma(t) = [a(t), b(t)]$, it follows from Theorem 1 that L_R^1 is closed (it is even $\sigma(L^1, L^\infty)$ compact!) whatever a and b are. The best idea is to consider the multifunction $t \mapsto \text{dom } \delta^*(\cdot | \Gamma(t))$. That will be stated precisely in Corollary 4.

5. GEOMETRIC RESULTS

The following result is an easy consequence of von Neumann–Aumann–Sainte–Beuve theorems [2, III.22, p. 74, III.23, p. 75].

LEMMA 2. If Γ and Γ' are μ -measurable multifunctions such that $L_R^1 = L_R^1 \upharpoonright_{\Gamma} \neq \phi$ then $\Gamma(t) = \Gamma'(t)$ μ -a.e.

THEOREM 3. (1) A necessary condition for L_R^1 to be closed in L_E^1 , is: $\Gamma(t)$ is a.e. closed and convex, and As $\Gamma(t) = [\mathcal{H}(t)]^0$ a.e.

(2) Suppose that $\dim E < \infty$ or Γ is LCNL valued. Then a sufficient condition for L_R^1 to be closed is:

$$\Gamma(t) \quad \text{is a.e. closed and convex, and} \quad \text{As } \Gamma(t) \supset [\mathcal{H}(t)]^0 \text{ a.e.}$$

(3) Suppose that $\dim E < \infty$ or $\Gamma(t)$ is contained in a fixed LCNL subset of E' . Then $\overline{L_R^{-1}} = L_E^{-1}$ with

$$\Sigma(t) = \overline{\text{co } \Gamma(t) + [\mathcal{H}(t)]^0}.$$

Proof. (1) By Theorem 0 $L_R^{-1} = L_E^{-1}$, and by Lemma 2 $\Gamma(t) = \Sigma(t)$ a.e. We noticed in Remark 2 that $C(t) = \text{As } \Sigma(t)$ a.e. Then $C(t) = \text{As } \Gamma(t)$ a.e.

(2) Let (φ_n) be the sequence of Theorem 0. Let (ψ_k) be the countably family of all $\sum_{i=0}^p \alpha_i \varphi_i$ for $p \in \mathbf{N}$, $\alpha_i \in \mathbf{Q}_+$. As \mathcal{H} is conic, one has $\psi_k \in \mathcal{H}$. Then one has

$$\begin{aligned} C(t) &= \{x' \mid \forall n, \langle x', \varphi_n(t) \rangle \leq 0\} \\ &= \{x' \mid \forall k, \langle x', \psi_k(t) \rangle \leq 0\}. \end{aligned}$$

As $\{\overline{\psi_k(t)} \mid k \in \mathbf{N}\}$ is conic we have

$$C(t) = [\overline{\{\psi_k(t) \mid k \in \mathbf{N}\}}]^0 \quad \text{a.e.} \quad (2)$$

Moreover a.e.

$$\begin{aligned} \text{As } \Gamma(t) &\supset [\mathcal{H}(t)]^0 \quad (\text{by hypothesis}) \\ &= C(t) \\ &\supset \text{As } \Gamma(t) \quad (\text{because } \Sigma \supset \Gamma) \end{aligned}$$

hence

$$C(t) = \text{As } \Gamma(t) \quad \text{a.e.} \quad (3)$$

From [2, I.7, p. 8]

$$\text{As } \Gamma(t) = [\text{dom } \delta^*(\cdot \mid \Gamma(t))]^0. \quad (4)$$

Combining (2), (3), (4) we obtain

$$[\text{dom } \delta^*(\cdot \mid \Gamma(t))]^0 = [\overline{\{\psi_k(t) \mid k \in \mathbf{N}\}}]^0 \quad \text{a.e.}$$

hence

$$\overline{\text{dom } \delta^*(\cdot \mid \Gamma(t))} = \overline{\{\psi_k(t) \mid k \in \mathbf{N}\}} \quad \text{a.e.} \quad (5)$$

We set $\Theta(t) = \{x' \mid \forall k, \langle x', \psi_k(t) \rangle \leq \delta^*(\psi_k(t) \mid \Gamma(t))\}$. By Theorem 1 L_{Θ}^{-1} is closed. Moreover it is obvious that $\Gamma(t) \subset \Theta(t) \subset \Sigma(t)$ a.e. hence $\overline{L_R^{-1}} = L_{\Theta}^{-1}$. It remains to show that $\Gamma(t) = \Theta(t)$ a.e.

(a) First suppose that $\Gamma(t)$ is LCNL. By [2, I.15] $\text{dom } \delta^*(\cdot \mid \Gamma(t))$ has a nonempty interior. So by (5) the $\psi_k(t)$ which belong to $\text{dom } \delta^*(\cdot \mid \Gamma(t))$ are dense in it. And by [2, III.34, p. 82], $\Gamma(t) = \Theta(t)$.

(b) Suppose now $\dim E < \infty$. The two cones in (5) are contained in the subspace generated by $\text{dom } \delta^*(\cdot \mid \Gamma(t))$. Denote by $A(t)$ the relative interior of $\text{dom } \delta^*(\cdot \mid \Gamma(t))$. By (5) the $\psi_k(t)$ which belong to $A(t)$ are dense in it. By

Rockafellar [10, Theorem 10.1, p. 82] $\delta^*(\cdot | \Gamma(t))$ is continuous on $A(t)$. So the arguments of [2, III.33 and III.34] remain valid and $\Gamma(t) = \Theta(t)$ a.e.

(3) First we have to prove that

$$\Sigma(t) = \overline{\text{co } \Gamma(t) + [\mathcal{H}(t)]^0}$$

is measurable. It follows from Theorem 0 that $t \mapsto [\mathcal{H}(t)]^0 = C(t)$ has a closed graph in $T \times E'_c$, where E'_c denotes E' endowed with the topology of uniform convergence on compact subsets of E . It is well known that E'_c is a Lusin space, so $\mathcal{B}(E'_c) = \mathcal{B}(E'_c)$. Hence C is measurable. From dense sequences of measurable selections of Γ and C it is easy (as in [2, III.40]) to deduce a dense sequence of measurable selections of Σ . So if $\dim E < \infty$, Σ is measurable by [2, III.30]. If K is a closed convex LCNL subset of E' such that $\Gamma(t) \subset K$ a.e. then \mathcal{H} contains the constant functions with values in $\text{dom } \delta^*(\cdot | K)$ and so $[\mathcal{H}(t)]^0 \subset [\text{dom } \delta^*(\cdot | K)]^0 = \text{As } K$. Therefore $\Sigma(t) \subset K$. Then the measurability of Σ follows from [2, III.37, p. 84]. Denote $\mathcal{H}_\Sigma = \text{dom } \delta^*(\cdot | L_\Sigma^1)$. From $\Sigma(t) \supset \Gamma(t)$ follows $\mathcal{H}_\Sigma \subset \mathcal{H}$. Moreover

$$\delta^*(\varphi(t) | \Sigma(t)) = \delta^*(\varphi(t) | \Gamma(t)) + \begin{cases} 0 & \text{if } \varphi(t) \in \overline{\mathcal{H}(t)} \\ +\infty & \text{if } \varphi(t) \notin \overline{\mathcal{H}(t)}. \end{cases}$$

So $\varphi \in \mathcal{H} \Rightarrow \varphi \in \mathcal{H}_\Sigma$. Finally $\mathcal{H} = \mathcal{H}_\Sigma$. Hence $\Sigma(t) \supset [\mathcal{H}(t)]^0 = [\mathcal{H}_\Sigma(t)]^0$. The second part of the present theorem ensures that L_Σ^1 is closed. Let us denote by Σ_0 the multifunction such that $\overline{L_\Gamma^1} = L_{\Sigma_0}^1$. It is clear that $\Sigma_0(t) \supset \text{co } \Gamma(t)$ a.e., and by Theorem 0 and Remark 2, $\text{As } \Sigma_0(t) = [\mathcal{H}(t)]^0$ a.e. hence $\Sigma_0(t) \supset \Sigma(t)$ a.e. From $L_\Gamma^1 \subset L_\Sigma^1 \subset L_{\Sigma_0}^1 = \overline{L_\Gamma^1}$ and the closedness of L_Σ^1 it follows $\overline{L_\Gamma^1} = L_\Sigma^1$.

Application of Theorem 3 to the Examples

EXAMPLE 1. One has for $\varphi \in \mathcal{C}$

$$\delta^*(\varphi | L_\Gamma^1) = \int_{[0,1]} \delta^*\left(\varphi(t) \left| \left[0, \frac{1}{t}\right]\right.\right) dt.$$

It is clear that $\varphi(0) > 0 \Rightarrow \delta^*(\varphi | L_\Gamma^1) = +\infty$ and that $\varphi(0) < 0 \Rightarrow \delta^*(\varphi | L_\Gamma^1) < \infty$. Then $\{\varphi | \varphi(0) < 0\} \subset \mathcal{H} \subset \{\varphi | \varphi(0) \leq 0\}$. So $]-\infty, 0[\subset \mathcal{H}(0) \subset]-\infty, 0[$ and $\forall t > 0$, $\mathcal{H}(t) = \mathbf{R}$, hence $[\mathcal{H}(0)]^0 = [0, \infty[$ and $\forall t > 0 [\mathcal{H}(t)]^0 = \{0\}$.

It follows from part (3) of Theorem 3 that

$$\begin{aligned} \Sigma(t) &= [0, \infty[& \text{if } t &= 0 \\ &= [0, 1/t] & \text{if } t > 0. \end{aligned}$$

EXAMPLE 2. Here if $\varphi \leq 0$, $\delta^*(\varphi | L_\Gamma^1) < \infty$. Otherwise there exists t such that $\varphi(t) > 0$ but as $T - K$ is open and dense in T one has

$\int \delta^*(\varphi(t) | \Gamma(t)) dt = +\infty$. So $\mathcal{H} = -\mathcal{C}_+$, hence $\forall t, \mathcal{H}(t) =]-\infty, 0]$ and $\Sigma(t) = [0, \infty[$.

Now we can develop the idea of Remark 5. Consider the two conditions

(C) There exists a closed convex-valued multifunction Γ' such that $\Gamma''(t) = \Gamma(t)$ a.e. and $\forall t, \overline{\mathcal{H}(t)} \supset \overline{\text{dom } \delta^*(\cdot | \Gamma'(t))}$.

(C') There exists a closed convex-conical-valued multifunction Γ' such that $\Gamma''(t) = \Gamma(t)$ a.e. and the multifunction $t \rightarrow [\Gamma'(t)]^0$ is l.s.c.

COROLLARY 4. (1) (C) is a necessary condition for L_{Γ^1} to be closed. Moreover if L_{Γ^1} is closed one can choose Γ' in (C) such that $\forall t, \overline{\mathcal{H}(t)} = \overline{\text{dom } \delta^*(\cdot | \Gamma'(t))}$.

(2) Suppose $\dim E < \infty$ or $\Gamma(t)$ is a.e. LCNL. Then (C) is a sufficient condition for L_{Γ^1} to be closed.

(3) (C') is a sufficient condition for L_{Γ^1} to be closed.

Remark 6. The relation $\overline{\mathcal{H}(t)} = \overline{\text{dom } \delta^*(\cdot | \Gamma'(t))}$ implies that $t \mapsto \overline{\text{dom } \delta^*(\cdot | \Gamma'(t))}$ is l.s.c. Indeed if U is an open set then

$$\begin{aligned} \{t | \overline{\mathcal{H}(t)} \cap U \neq \emptyset\} &= \{t | \mathcal{H}(t) \cap U \neq \emptyset\} \\ &= \cup \{\varphi^{-1}(U) | \varphi \in \mathcal{H}\} \end{aligned}$$

is open. But if Γ is not conical valued, the l.s. continuity of $t \mapsto \overline{\text{dom } \delta^*(\cdot | \Gamma'(t))}$ is not sufficient to ensure L_{Γ^1} to be closed. Indeed in Example 1, $\forall t, \text{dom } \delta^*(\cdot | \Gamma(t)) = \mathbf{R}$.

Remark 7. Rockafellar ([11, Corollary 5B, p. 466]) proves that L_{Γ^1} is closed (and also that $\forall t, \text{As } \Gamma(t) = C(t)$) under the following hypotheses:

(a) $\forall t, \Gamma(t)$ is LCNL.

(b) $t \mapsto \text{dom } \delta^*(\cdot | \Gamma(t))$ is fully l.s.c. (that is a stronger assumption than l.s.c.-ness).

(c) $\forall V$ open subset of $T, \forall x \in E = \mathbf{R}^n$ such that \exists a neighborhood U of x such that $\forall t \in V, \text{dom } \delta^*(\cdot | \Gamma(t)) \supset U$, then

$$\int_V |\delta^*(x | \Gamma(t))| \mu(dt) < \infty.$$

(d) the support of μ is T .

Proof. (1) By Theorem 3(1), As $\Gamma(t) = [\mathcal{H}(t)]^0$ a.e. By [2, I.7, p. 8] As $\Gamma(t) = [\text{dom } \delta^*(\cdot | \Gamma(t))]^0$ hence $\overline{\text{dom } \delta^*(\cdot | \Gamma(t))} = \overline{\mathcal{H}(t)}$ a.e. We choose

$$\begin{aligned} \Gamma'(t) &= \Gamma(t) & \text{if } & \text{As } \Gamma(t) = [\mathcal{H}(t)]^0 \\ &= [\mathcal{H}(t)]^0 & \text{otherwise.} \end{aligned}$$

So we have $\forall t, \overline{\text{dom } \delta^*(\cdot \mid \Gamma'(t))} = \overline{\mathcal{H}(t)}$. That is clear in the first eventuality, and in the second it follows from the fact that $\mathcal{H}(t)$ is conic.

(2) One has

$$\text{As } \Gamma'(t) = [\text{dom } \delta^*(\cdot \mid \Gamma'(t))]^0 \supset [\mathcal{H}(t)]^0.$$

So by Theorem 3(2), L_{r^1} is closed.

(3) Let \mathcal{H} denote the set of continuous selections of $t \mapsto [\Gamma'(t)]^0$. As T is compact and E Fréchet, the Michael theorem [5] ensures $\mathcal{H}(t) = [\Gamma'(t)]^0$. Let (φ_n) be a dense sequence in \mathcal{H} (for the uniform convergence topology). Then

$$\forall t, \overline{\{\varphi_n(t) \mid n \in \mathbf{N}\}} = [\Gamma'(t)]^0.$$

So one has

$$\Gamma'(t) = \{x' \mid \forall n, \langle x', \varphi_n(t) \rangle \leq 0\},$$

and by Theorem 1, L_{r^1} is closed.

6. FUNCTIONAL RESULTS

We set again some notation and hypotheses.

Let $f: T \times E' \rightarrow \bar{\mathbf{R}}$ be an integrand, that is a measurable function. We shall write f_t for $f(t, \cdot)$. We define $I_f(u)$ for $u \in \mathcal{L}_{E'}^1$ by

$$I_f(u) = \int f(t, u(t)) \mu(dt)$$

with the convention that if both the negative and positive parts of $f(t, u(t))$ are nonintegrable, then $I_f(u) = +\infty$.

To f corresponds the multifunction $\Gamma(t) = \text{epi } f_t$ with values in $E' \times \mathbf{R}$. By [2, VII.1, p. 196] Γ is measurable. The hypotheses (H1) and (H2) relative to Γ become

(H'1) μ is nonatomic or f_t is a.e. convex.

(H'2) $\exists u \in L_{E'}^1$ such that $I_f(u) < \infty$ ($I_f(u) = -\infty$ is allowed).

Indeed

$$(H2) \Leftrightarrow \exists (u, v) \in L_{r^1}$$

$$\Leftrightarrow \exists u \in L_{E'}^1, \quad \exists v \in L_{\mathbf{R}}^1 \quad \text{such that} \quad f(t, u(t)) \leq v(t) \quad \text{a.e.}$$

$$\Leftrightarrow (H'2).$$

Under these hypotheses we associate to I' the set

$$\mathcal{H} = \text{dom } \delta^*(\cdot \mid L_{I'}^1) = \{(\varphi, \alpha) \in \mathcal{C}_{E \times \mathbb{R}} \mid \delta^*(\varphi, \alpha) \mid L_{I'}^1 < \infty\},$$

and the multifunctions Σ and C defined in Theorem 0.

It follows from $\Sigma(t) \supset I'(t)$ a.e. that $\Sigma(t)$ is a.e. the epigraph of a convex l.s.c. function g_t . By [2, VII.1, p. 196] the function $g: T \times E' \rightarrow \bar{\mathbf{R}}$ defined by $g(t, x') = g_t(x')$ is an integrand. It will be proved in Lemma 8 that $C(t)$ is also an epigraph. Consider $f^*: T \times E \rightarrow \bar{\mathbf{R}}$ defined by $f^*(t, \cdot) = [f(t, \cdot)]^*$. We define $I_{f^*}(\varphi)$ for $\varphi \in \mathcal{C}_E$ by

$$I_{f^*}(\varphi) = \int f^*(t, \varphi(t)) \mu(dt).$$

Thanks to a famous theorem of Rockafellar [2, VII.7, p. 200], under (H'2)

$$I_{f^*}(\varphi) = (I_f)^*(\varphi) = \sup\{\langle u, \varphi \rangle - I_f(u) \mid u \in L_{E'}^1\}.$$

Let

$$\mathcal{G} = \text{dom } I_{f^*} = \{\varphi \in \mathcal{C}_E \mid I_{f^*}(\varphi) < \infty\}.$$

We assume the following hypothesis

$$(H3) \quad \mathcal{G} \neq \emptyset.$$

That means $\exists \varphi_0 \in \mathcal{C}_E$ such that $\int f^*(t, \varphi_0(t)) \mu(dt) < \infty \Leftrightarrow \exists \varphi_0 \in \mathcal{C}_E$ $\exists \psi \in \mathcal{L}_{\mathbb{R}}^1$ such that $f^*(t, \varphi_0(t)) \leq \psi(t)$ a.e. $\Leftrightarrow \exists \varphi_0 \in \mathcal{C}_E, \exists \psi \in \mathcal{L}_{\mathbb{R}}^1$ such that for almost every t ,

$$\forall x', \langle x', \varphi_0(t) \rangle - f(t, x') \leq \psi(t).$$

So (H3) is equivalent to

$$(H'3) \quad \exists \varphi_0 \in \mathcal{C}_E, \exists \psi \in \mathcal{L}_{\mathbb{R}}^1 \text{ such that for almost every } t,$$

$$\forall x', f(t, x') \geq \langle x', \varphi_0(t) \rangle - \psi(t).$$

Remark 8. If I_f is a convex proper $\sigma(L_{E'}^1, \mathcal{C}_E)$ l.s.c. functional, then its polar I_{f^*} is also proper, so (H3) is satisfied. Moreover under (H'1) it will be proved in Lemma 6 that if I_f is l.s.c. it is necessarily convex.

The following hypothesis is assumed on and after Lemma 8:

$$(H4) \quad \text{The support of } \mu \text{ is } T.$$

LEMMA 5. *Let f be an integrand. Let $A: L_{E' \times \mathbb{R}}^1 \rightarrow L_{E'}^1 \times \mathbf{R}$ be defined by $A(u, v) = (u, \int v d\mu)$. Then $\text{epi } I_f = A(L_{I'}^1)$.*

Proof. Let $u \in \mathcal{L}_E^1$, $r \in \mathbf{R}$. Then

$$\begin{aligned} (u, r) \in \text{epi } I_f &\Leftrightarrow r \geq I_f(u) \\ &\Leftrightarrow \exists v \in \mathcal{L}_R^1 \end{aligned}$$

such that

$$v(t) \geq f(t, u(t)) \quad \text{a.e.} \quad \text{and} \quad \int v \, d\mu = r$$

(we prove \Rightarrow below)

$$\Leftrightarrow \exists v \in \mathcal{L}_R^1 \quad \text{such that} \quad (u, v) \in \mathcal{L}_R^1 \quad \text{and} \quad A(u, v) = (u, r).$$

We prove now the second \Rightarrow . Thanks to the convention in the definition of I_f ,

$$r \geq I_f(u) \Rightarrow f(t, u(t))^+ \text{ is integrable.}$$

If $f(t, u(t))^-$ is also integrable we put

$$v(t) = f(t, u(t)) + \frac{r - I_f(u)}{\mu(T)}.$$

If $f(t, u(t))^-$ is not integrable, then for every $n \geq 0$ $h_n(t) = \sup(-n, f(t, u(t)))$ is integrable. By the Fatou lemma, for n large enough, one has $\int h_n \leq r$. So we put

$$v = h_n + \frac{r - \int h_n \, d\mu}{\mu(T)}. \quad \blacksquare$$

From now on we assume (H'1), (H'2), and (H3).

LEMMA 6. *The l.s.c. hull of I_f is convex and its epigraph is $\overline{\text{epi } I_f}$. The polar of I_{f*} within the duality $\mathcal{C}_E, \mathcal{M}_{E'}$ is convex and its epigraph is $\overline{\text{epi } I_f}$.*

Proof. As A is continuous, the formula

$$\overline{\text{epi } I_f} = \overline{A(L_R^1)} \quad \text{implies} \quad \overline{\text{epi } I_f} = \overline{A(\overline{L_R^1})}.$$

The Liapunov theorem implies that $\overline{L_R^1}$ is convex. So the first assertion is proved.

(2) Let $\hat{I}_f: \mathcal{M}_{E'} \rightarrow \bar{\mathbf{R}}$ be defined by

$$\begin{aligned} \hat{I}_f(m) &= I_f(u) & \text{if} & \quad m = u \cdot \mu \\ &= +\infty & \text{if} & \quad m \text{ is not absolutely continuous with respect to } \mu. \end{aligned}$$

We said above that $I_{f*} = (I_f)^*$. One has also

$$I_{f*} = (\hat{I}_f)^* \quad \text{within the duality} \quad \mathcal{M}_{E'}, \mathcal{C}_E.$$

Moreover \hat{I}_f is greater than the following continuous affine function

$$m \mapsto \langle m, \varphi_0 \rangle - \int \psi \, d\mu \quad (\text{thanks to } (H'3)).$$

So by a well known result [2, I.5, p. 6] the greatest convex l.s.c. function less than \hat{I}_f is its bipolar $(\hat{I}_f)^{**} = (I_f^*)^*$. Hence

$$\begin{aligned} \text{epi}(I_f^*)^* &= \overline{\overline{\text{co epi } \hat{I}_f}} = \overline{\overline{\text{co epi } I_f}} \\ &= \overline{\overline{\text{epi } I_f}} \quad \text{since} \quad \overline{\overline{\text{epi } I_f}} \text{ is convex.} \quad \blacksquare \end{aligned}$$

We need now the adjoints A^* and A^{**} . They are easy to calculate:

$$A^*: \mathcal{C}_E \times \mathbf{R} \rightarrow \mathcal{C}_{E \times \mathbf{R}} = \mathcal{C}_E \times \mathcal{C}_{\mathbf{R}}$$

satisfies

$$A^*(\varphi, r) = (\varphi, r\chi_T),$$

and

$$A^{**}: \mathcal{M}_{E'} \times \mathcal{M}_{\mathbf{R}} \rightarrow \mathcal{M}_{E'} \times \mathbf{R}$$

satisfies

$$A^{**}(\lambda, \sigma) = (\lambda, \sigma(T)).$$

LEMMA 7. One has $\overline{\overline{\text{epi } I_f}} = A^{**}(\overline{\overline{L_T^1}})$.

Proof. From Lemma 5 and the continuity of A^{**} follows $\overline{\overline{\text{epi } I_f}} \supset A^{**}(\overline{\overline{L_T^1}})$. Conversely let $(\lambda, r) \in \overline{\overline{\text{epi } I_f}}$.

There exists a generalized sequence $(u_i, r_i)_{i \in I}$ in $\text{epi } I_f$ which converges to (λ, r) . There exists $v_i \in L_{\mathbf{R}}^1$ such that $(u_i, v_i) \in L_T^1$ and $\int v_i \, d\mu = r_i$.

By (H'3) $f(t, x') \geq \langle x', \varphi_0(t) \rangle - \psi(t)$. So

$$-v_i(t) + \langle u_i(t), \varphi_0(t) \rangle \leq -f(t, u_i(t)) + \langle u_i(t), \varphi_0(t) \rangle \leq \psi(t).$$

One has

$$\begin{aligned} \int (-v_i + \langle u_i, \varphi_0 \rangle)^- \, d\mu &= \int (-v_i + \langle u_i, \varphi_0 \rangle)^+ \, d\mu - \int (-v_i + \langle u_i, \varphi_0 \rangle) \, d\mu \\ &\leq \int \psi^+ \, d\mu - \int (-v_i + \langle u_i, \varphi_0 \rangle) \, d\mu \end{aligned}$$

and

$$\int (-v_i + \langle u_i, \varphi_0 \rangle) \, d\mu \rightarrow -r + \langle \lambda, \varphi_0 \rangle.$$

Hence for i large enough, say $i \geq i_0$ $\int (-v_i + \langle u_i, \varphi_0 \rangle)^- \, d\mu$ is bounded and so $\int |-v_i + \langle u_i, \varphi_0 \rangle| \, d\mu$ is also bounded.

The real measures $(-v_i + \langle u_i, \varphi_0 \rangle) \mu$ for $i \geq i_0$ belongs to a weakly compact subset of $\mathcal{M}_{\mathbb{R}}$. Let \mathcal{U} be an ultrafilter on I finer than the filter of sections, the limit $\lim_{\mathcal{U}} (-v_i + \langle u_i, \varphi_0 \rangle) \mu$ exists. It is easy to see that $\langle u_i, \varphi_0 \rangle \mu$ converges to the real Radon measure $\alpha \mapsto \langle \lambda, \alpha \varphi_0 \rangle$, hence $\lim_{\mathcal{U}} v_i \mu$ exists. Thanks to the continuity of A^{**} we have

$$(\lambda, r) = A^{**}(\lambda, \lim_{\mathcal{U}} v_i \mu),$$

where $(\lambda, \lim_{\mathcal{U}} v_i \mu) = \lim_{\mathcal{U}} (u_i, v_i)$ belongs to $\overline{L_R^1}$.

LEMMA 8. For every t , $C(t) = \text{epi } \delta^*(\cdot \mid \mathcal{G}(t))$ (where $\mathcal{G}(t) = \{\varphi(t) \mid \varphi \in \mathcal{G}\}$).

Proof. We have to prove $[\mathcal{H}(t)]^0 = \text{epi } \delta^*(\cdot \mid \mathcal{G}(t))$. First remark that

$$\begin{aligned} \varphi \in \mathcal{G} &\Leftrightarrow \int f^*(t, \varphi(t)) \mu(dt) < \infty \\ &\Leftrightarrow \int \delta^*((\varphi(t), -1) \mid \text{epi } f_t) \mu(dt) < \infty \\ &\Leftrightarrow \delta^*((\varphi, -\chi_T) \mid L_{T^1}) < \infty \\ &\Leftrightarrow (\varphi, -\chi_T) \in \mathcal{H}. \end{aligned}$$

If $(x', r) \in [\mathcal{H}(t)]^0$, for every $\varphi \in \mathcal{G}$, one has $\langle (x', r), (\varphi(t), -1) \rangle \leq 0$ hence $r \geq \langle x', \varphi(t) \rangle$ which implies $(x', r) \in \text{epi } \delta^*(\cdot \mid \mathcal{G}(t))$. It remains to prove $\text{epi } \delta^*(\cdot \mid \mathcal{G}(t)) \subset [\mathcal{H}(t)]^0$. Let $(x', r) \in \text{epi } \delta^*(\cdot \mid \mathcal{G}(t))$ and $(\varphi, \alpha) \in \mathcal{H}$. We have to prove $\langle (x', r), (\varphi(t), \alpha(t)) \rangle \leq 0$.

First let us prove $\forall s, \alpha(s) \leq 0$. One has

$$\delta^*((\varphi, \alpha) \mid L_{T^1}) = \int \delta^*((\varphi(s), \alpha(s)) \mid \text{epi } f_s) \mu(ds) < \infty.$$

As $\text{epi } f_s \neq \emptyset$ a.e., $\alpha(s)$ is ≤ 0 a.e. But α is a continuous function and the support of μ is T , so $\forall s, \alpha(s) \leq 0$.

(a) The simplest case is when $\forall s, \alpha(s) < 0$. Therefore $-1/\alpha$ is a positive continuous, hence bounded function. So

$$\begin{aligned} \delta^*\left(-\frac{1}{\alpha}(\varphi, \alpha) \mid L_{T^1}\right) &= \int \delta^*\left(-\frac{1}{\alpha}(\varphi, \alpha) \mid \Gamma(\cdot)\right) d\mu \\ &= \int -\frac{1}{\alpha} \delta^*((\varphi, \alpha) \mid \Gamma(\cdot)) d\mu \end{aligned}$$

is finite.

Hence $-(1/\alpha)(\varphi, \alpha) = (-\varphi/\alpha, -\chi_T)$ belongs to \mathcal{H} , and $-\varphi/\alpha \in \mathcal{G}$. Therefore $r \geq \langle x', -\varphi(t)/\alpha(t) \rangle$ or $\langle x', \varphi(t) \rangle + r\alpha(t) \leq 0$.

(b) Consider now the general case. Let $\varphi_0 \in \mathcal{G}$. For every $\lambda \in [0, 1[$ $(\psi, \beta) = \lambda(\varphi, \alpha) + (1 - \lambda)(\varphi_0, -\chi_T)$ belongs to \mathcal{H} and $\forall s, \beta(s) < 0$. So by (a)

$$\langle x', \psi(t) \rangle + r\beta(t) \leq 0.$$

As λ can converge to 1, one has $\langle x', \varphi(t) \rangle + r\alpha(t) \leq 0$. ■

The Lebesgue decomposition theorem is valid in $\mathcal{M}_{E'}$ ([13, Remark 2 after Theorem 1]). If $m \in \mathcal{M}_{E'}$ one has $m = (dm/d\mu)\mu + m_s$, where m_s is μ -singular. There exists a real bounded positive measure θ such that m_s is absolutely continuous with respect to θ . Then $m = (dm/d\mu) + (dm_s/d\theta)\theta$.

THEOREM 9. *With the foregoing notation one has*

$$(I_{f*})^*(m) = I_g\left(\frac{dm}{d\mu}\right) + \int \delta^*\left(\frac{dm_s}{d\theta}(t) \mid \mathcal{G}(t)\right) \theta(dt). \quad (6)$$

Remark 9. The last term does not depend on the measure θ .

Proof. (1) First we show that the right-hand side is meaningful. We have noticed in the proof of part 3 of Theorem 3 that C is measurable, then by Lemma 8 and [2, VIII.1, p. 196], $(t, x') \mapsto \delta^*(x' \mid \mathcal{G}(t))$ is an integrand.

Let $\varphi_0 \in \mathcal{G}$, then

$$\int \delta^*\left(\frac{dms}{d\theta}(t) \mid \mathcal{G}(t)\right) \theta(dt) \geq \int \left\langle \frac{dms}{d\theta}, \varphi_0 \right\rangle d\theta > -\infty.$$

Let φ_0 and ψ satisfying (H'3). Consider

$$\Gamma_1(t) = \text{epi}(x' \mapsto \langle x', \varphi_0(t) \rangle - \psi(t)).$$

By Theorem 1, $L_{\Gamma_1}^1$ is closed, so that $\Sigma(t) \subset \Gamma_1(t)$ a.e., that is,

$$g(t, x') \geq \langle x', \varphi_0(t) \rangle - \psi(t).$$

Therefore $I_g(dm/d\mu) > -\infty$

(2) By Lemma 7

$$(m, r) \in \overline{\text{epi } I_f} \Leftrightarrow \exists (m, \sigma) \in \overline{L_R^1} \quad \text{such that} \quad \sigma(T) = r.$$

By Lemma 6

$$\begin{aligned} (I_{f*})^*(m) &= \inf\{r \mid (m, r) \in \overline{\text{epi } I_f}\} \\ &= \inf\{\sigma(T) \mid (m, \sigma) \in \overline{L_R^1}\}. \end{aligned}$$

Let $(m, \sigma) \in \overline{L_R^1}$. We shall prove

$$\sigma(T) \geq I_g \left(\frac{dm}{d\mu} \right) + \int \delta^* \left(\frac{dm_s}{d\theta}(t) \mid \mathcal{G}(t) \right) \theta(dt). \quad (7)$$

Let (m_s, σ_s) be the μ -singular part of (m, σ) . By Theorem 0 there exists a real positive measure θ singular with respect to μ such that (m_s, σ_s) is absolutely continuous with respect to θ and

$$\left(\frac{dm_s}{d\theta}, \frac{d\sigma_s}{d\theta} \right) \in L_{\mathcal{C}}^1(\theta), \quad (8)$$

$$\left(\frac{dm}{d\mu}, \frac{d\sigma}{d\mu} \right) \in L_{\mathcal{E}}^1. \quad (9)$$

From (8) follows $(d\sigma_s/d\theta)(t) \geq \delta^*((dm_s/d\theta)(t) \mid \mathcal{G}(t)) \theta$ -a.e. so that $\sigma_s(T) \geq \int \delta^*((dm_s/d\theta)(t) \mid \mathcal{G}(t)) \theta(dt)$.

From (9) follows $(d\sigma/d\mu)(t) \geq g(t, (dm/d\mu)(t)) \mu$ a.e. so that $\int (d\sigma/d\mu) d\mu \geq I_g(dm/d\mu)$. As $\sigma(T) = \int (d\sigma/d\mu) d\mu + \sigma_s(T)$, inequality (7) is proved.

(3) To prove (6) it remains to show that if the right-hand side is finite, there exists σ such that $(m, \sigma) \in \overline{L_R^1}$ and (7) becomes an equality. That is true because it suffices to put

$$\sigma = g \left(\cdot, \frac{dm}{d\mu}(\cdot) \right) \mu + \delta^* \left(\frac{dm_s}{d\theta}(\cdot) \mid \mathcal{G}(\cdot) \right) \theta.$$

COROLLARY 10. *The l.s.c. hull of I_f is I_g . If $\dim E < \infty$ or if $\text{epi } f_t$ is contained in a fixed closed convex LCNL subset of $E' \times \mathbf{R}$ one has*

$$\text{epi } g_t = \overline{\text{co epi } f_t + \text{epi } \delta^*(\cdot \mid \mathcal{G}(t))}.$$

Proof. By Lemma 6 the l.s.c. hull of I_f is equal to $(I_{f*})^*$ restricted to L_E^1 . By Theorem 9 it is I_g . The second assertion follows from Theorem 3 (3).

COROLLARY 11. *Suppose $\dim E < \infty$ or $\text{epi } f_t$ is a.e. LCNL. The following condition is sufficient for I_f to be l.s.c.: f_t is a.e. convex l.s.c. and $\text{rec } f_t \leq \delta^*(\cdot \mid \mathcal{G}(t))$ a.e. (where $\text{rec } f_t$ is defined by $\text{epi}(\text{rec } f_t) = \text{As}(\text{epi } f_t)$).*

Proof. The condition $\text{rec } f_t \leq \delta^*(\cdot \mid \mathcal{G}(t))$ means $\text{As } \Gamma(t) \supset C(t)$. So by Theorem 3(2), L_R^1 is closed, hence $\Gamma(t) = \Sigma(t)$ a.e. that is, $f_t = g_t$ a.e. By Corollary 10 I_f is l.s.c.

COROLLARY 12. *The following condition is necessary for I_f to be l.s.c.: f_t is a.e. convex l.s.c. and $\text{rec } f_t = \delta^*(\cdot \mid \mathcal{G}(t))$ a.e. (where $\text{rec } f_t$ is defined by $\text{epi}(\text{rec } f_t) = \text{As}(\text{epi } f_t)$).*

Proof. If I_f is l.s.c. it follows from Corollary 10 that $I_f = I_g$. Thanks to the next lemma, $f_t = g_t$ a.e. So $L_f^1 = L_g^1$ is closed, and the necessary condition follows from Theorem 3(1).

LEMMA 13. *Let f and g be two integrands such that:*

- (a) $g_t \leq f_t$ a.e.
- (b) $\exists u \in L_E^1$, such that $I_f(u) < \infty$,
- (c) $\forall u \in L_E^1$, $f(t, u(t))^-$ is integrable,
- (d) $I_f = I_g$.

Then $f_t = g_t$ a.e.

Proof. Let $\Gamma(t) = \text{epi } f_t$, $\Sigma(t) = \text{epi } g_t$. By (a) $\Sigma(t) \supset \Gamma(t)$ a.e. By (b) $L_f^1 \neq \emptyset$. By Lemma 2 if $f_t = g_t$ a.e. is not true one has $L_f^1 \neq L_g^1$. So there exists $(u, v) \in L_E^1 - L_f^1$.

Put $v'(t) = \inf(v(t), f(t, u(t)))$.

Thanks to (c) v' is still integrable. The function v' has the following properties.

- (1) $g(t, u(t)) \leq v'(t) \leq f(t, u(t))$,
- (2) $v'(t) < f(t, u(t))$ on a nonnegligible set, because $(u, v) \notin L_f^1$.

So we have $I_g(u) \leq \int v' d\mu < I_f(u)$ which is a contradiction to (d).

Remark 10. In Olech [7], $E = \mathbf{R}^n$, so our Corollaries 11 and 12 contain his characterization: I_f is l.s.c. iff L_f^1 is closed.

EXAMPLE 3. Let $T = [0, 1]$, $\mu = dt + \delta_0$, $E = \mathbf{R}$

$$\begin{aligned} f_t &= x \mapsto x^2 & \text{if } t &= 0 \\ &= x \mapsto t^2 x^2 & \text{if } t > 0. \end{aligned}$$

Then I_f is not l.s.c. because if $u_n = n\chi_{[0, 1/n]}$ one has $u_n \rightarrow \chi_{\{0\}}$, but

$$I_f(u_n) = \int_0^{1/n} t^2 n^2 dt = \frac{1}{3n} \rightarrow 0 \quad \text{and} \quad I_f(\chi_{\{0\}}) = 1.$$

We can calculate g . Indeed

$$\begin{aligned} f_t^* &= x' \mapsto \frac{1}{4} x'^2 & \text{if } t &= 0 \\ &= x' \mapsto \frac{1}{4t^2} x'^2 & \text{if } t > 0. \end{aligned}$$

So if $\varphi \in \mathcal{C}_{\mathbf{R}}$ one has $I_{f^*}(\varphi) < \infty \Rightarrow \varphi(0) = 0$.

Hence $\mathcal{G}(0) = \{0\}$ and it is clear that $\mathcal{G}(t) = \mathbf{R}$ for $t > 0$. By Corollary 10

$$\begin{aligned} g_t &= 0 && \text{if } t = 0 \\ &= x \mapsto t^2 x^2 && \text{if } t > 0. \end{aligned}$$

REFERENCES

1. H. BRÉZIS, Intégrales convexes dans les espaces de Sobolev, *Israel J. Math.* **13** (1972) 9–23.
2. C. CASTAING AND M. VALADIER, "Convex Analysis and Measurable Multifunctions," Lecture in Mathematics No. 580, Springer-Verlag, Berlin, 1977.
3. A. D. IOFFE, Sur la semi-continuité des fonctionnelles intégrales, *C. R. Acad. Sci. Paris* **284** (1977), 807–809.
4. A. D. IOFFE, On lower semicontinuity of integral functionals, I, *SIAM J. Contr. Optimization* **15** (1977), 521–538.
5. E. MICHAEL, Continuous selections, I, *Ann. of Math.* **63**, No. 2 (1956), 361–382.
6. C. OLECH, The characterization of the weak* closure of certain sets of integrable functions, *SIAM J. Contr.* **12**, No. 2 (1974), 311–318.
7. C. OLECH, A necessary and sufficient condition for lower semicontinuity of certain integral functionals, in "Mathematical structures, Computational Mathematics, Mathematical Modelling," pp. 373–379, Sofia, 1975.
8. C. OLECH, Existence theory in optimal control problem: The underlying ideas, in "International Conference on Differential Equations," pp. 612–635, Academic Press, New York, 1975.
9. C. OLECH, Existence theory in optimal control, in "Control Theory and Topics in Functional Analysis," Vol. I, pp. 291–328, International Atomic Energy Agency, Vienna, 1976.
10. R. T. ROCKAFELLAR, "Convex Analysis," Princeton Univ. Press, Princeton, N. J., 1970.
11. R. T. ROCKAFELLAR, Integrals which are convex functionals, II, *Pacific J. Math.* **39** (1971), 439–469.
12. R. T. ROCKAFELLAR, Integral functionals, normal integrals and measurable selections, in "Nonlinear Operators and the Calculus of Variations" (J. P. Gossez, Ed.), pp. 157–207, Lecture Notes in Mathematics No. 543, Springer-Verlag, Berlin, 1976.
13. M.-F. SAINTE-BEUVE, Some topological properties of vector measures with bounded variations and its applications, *Ann. Mat. Pura Appl.* **116** (1978), 317–379.
14. M.-F. SAINTE-BEUVE, Sur l'adhérence de certains ensembles de fonctions vectorielles intégrables, *Sém. d'An. Convexe Montpellier* No. 6 (1974).
15. M.-F. SAINTE-BEUVE, Propriétés topologiques des mesures vectorielles et quelques applications, *Sém. d'An. Convexe Montpellier* No. 2 (1975).
16. M.-F. SAINTE-BEUVE, Complément sur les applications des propriétés topologiques des mesures vectorielles, *Sém. d'An. Convexe Montpellier* No. 5 (1975).
17. M.-F. NOUGUÈS-SAINTE-BEUVE, Semi-continuité inférieure de certaines fonctionnelles intégrales, *Sém. d'An. Convexe Montpellier* No. 7 (1976).
18. M.-F. SAINTE-BEUVE, Propriétés topologiques des mesures vectorielles et applications, *C. R. Acad. Sci. Paris* **280** (1975), 881–884.
19. M.-F. SAINTE-BEUVE, Mesures vectorielles à variation bornée et représentation intégrale, in "Intégration vectorielle et multivoque" (A. Costé, Ed.), Actes du Colloque de Caen, Offilib, Paris, 1975.

20. TRAN CAO NGUYEN, A characterization of some weak lower semicontinuity of integral functionals, to appear.
21. M. VALADIER, Fermeture, régularisée s.c.i. et bipolaire vagues, *C. R. Acad. Sci. Paris* **284** (1977), 1025–1027.
22. M. VALADIER, Fermeture étroite et bipolaire vague, *Sém. d'An. Convexe Montpellier* No. 6 (1977).
23. TRAN CAO NGUYEN, Decomposition of the conjugate integral functional on the space of regular measures, to appear.